Double Balloon Endoscopy Prep:

WHAT YOU NEED:

- One large bottle of **polyethylene glycol 3350**. The most common brand is MiraLAX® but ClearLAX, Powderlax, etc. are examples of generic store brands.
- One package of senna of your choice. Senna comes in tablets, chocolate squares, and tea.
 If you need a liquid formulation and your child is 10 years or older, ask the GI clinic for a prescription.











EXAMPLES OF POLYETHLENE GLYCOL 3350 PRODUCTS

EXAMPLES OF SENNA PRODUCTS

HOW TO USE MiraLAX®:

- MiraLAX is a stool softener for children ages 6 months and older. It works by bringing more water into the colon and softening the stool, making it easier to pass.
- MiraLAX comes in a powder that is mixed in clear liquid.
- One dose = 1 capful of powder + 8 ounces (1 cup) of one of the following clear liquids:
 - Water
 - Flavored water
 - Jell-O[®] gelatin
 - Slushies
 - Icee®

- Popsicles
- Gatorade®
- Powerade®
- Pedialyte®
- Kool-Aid®

- Sprite[®]
- Starry
- Apple juice
- White grape juice





AVOID:

- All solid foods
- Milk or dairy products
- Red- or purple-colored liquids
- Orange juice or juice with pulp
- Gatorade "Recovery" brand, which contains whey protein that should be avoided during prep

The goal of the bowel prep is clear stool from your child's intestine so we can safely see and navigate during the procedure.

- During the prep your child's stool should become light yellow to clear in color.
- It is important that your child stays hydrated while doing their prep. Encourage extra clear liquids!
- Poor bowel prep will result in cancelling or rescheduling the procedure.

NURSE'S LINES

Gastroenterology Clinic Red Team Nurse Line (816) 760-8851
Gastroenterology Clinic Blue Team Nurse Line (816) 760-8852
Gastroenterology Clinic Green Team Nurse Line (816) 760-8853
Gastroenterology Pink Team Nurse Line (816) 983-6975

Gastroenterology Liver Team Nurse Line (816) 302-3410 OPT.2



Copyright © 2024 The Children's Mercy Hospital. All rights reserved.25-GI-0287 09/24

PLEASE FOLLOW THIS SCHEDULE

At 5 p.m. (two days before procedure)

- Begin clear liquid diet (see list of acceptable clear liquids). No solid food should be eaten.
- Give extra clear liquids in between doses of MiraLAX. Your child can have as much clear liquid as they want – this is important to help keep your child hydrated.

□ 5 p.m.	□ 7 p.m.	
□ 6 p.m.	□ 8 p.m.	

Next: Give one dose of MiraLAX each hour as scheduled below

ur as scheduled below	
☐ 9 a.m. Also give 2 senna tablets if your child is older than 10 years old	☐ 2 p.m. Please call GI Clinic Nurse list below if your of has not yet postool.
□ 10 a.m.	□ 3 p.m.
□ 11 a.m.	
□ 12 p.m.	□ 4 p.m.
	□ 5 p.m.
□ lom	

hild ssed

If your child's stools are clear or light yellow, they are ready for their procedure.

If stools are not clear or light yellow continue giving more hourly doses of MiraLAX.

ADDITIONAL INFORMATION

If you have questions, please call Nurse's Line listed to left during clinic hours.7:00 a.m. - 4:00 p.m.

If you need Additional assistance after hours, please call 816-234-3000 and ask to speak with the GI Doctor on call for further recommendations.