





QR code for mobile view

Inclusion Criteria:

- Patients undergoing
 - Ganz Periacetabular Osteotomy
 - Surgical Hip Dislocation

Exclusion Criteria:

- Closed pelvic procedures




Medication and Diet Instructions
(received at Orthopaedic Surgery Clinic and PAT):

- **Medication:**
 - Mupirocin for 5 days
 - Patient takes all normal daily medications night prior to surgery unless specifically instructed to stop
- **Diet:**
 - Standard NPO guidelines
 - **2 - 3 hrs prior to surgery:** Carbohydrate-rich drink (Gatorade, Powerade, or Pedialyte)
- **Arrival time/location**

Patient seen in **Orthopaedic Surgery Clinic AND PAT** for pre-surgery evaluation to review the following:

- Pain management and regional anesthesia plan, including risks
- Medication and diet instructions
- Mitigation of pre-op anxiety
- Refer to PT for gait training
 - *Same day as pre-op appointment*
- [ERAS overview handout \(English\)](#)
- [ERAS overview handout \(Spanish\)](#)
- **Medications to be ordered in PAT**
 - Celecoxib (Patients >25 kg, 100 mg PO)
 - Midazolam as indicated



Abbreviations:

PAO = periacetabular osteotomy
PAT = pre-admission testing
SDS = same day surgery

[Intraoperative algorithm](#)
[Inpatient algorithm](#)